arXiv:1208.0511v1 [math.DS] 2 Aug 2012

BILLIARD DYNAMICS NEAR AN INVARIANT HORIZONTAL CIRCLE

GERALDO CÉSAR GONÇALVES FERREIRA, SYLVIE OLIFFSON KAMPHORST,
AND SÔNIA PINTO-DE-CARVALHO

Abstract. Sufficiently differentiable oval billiards always have invariant rotational curves, but there are only two types of ovals with an invariant horizontal circle in its phase-space: the constant width ovals and some very special symmetric curves.

In this work we study the dynamics near the horizontal circle for the billiard map of those two types of ovals, and show that the horizontal circle is approached, from both sides, by other invariant rotational curves. We will also describe some dynamical consequences.

1. Introduction

Let Γ be an oval: a plane, simple, closed curve, with strictly positive curvature. The billiard problem on Γ consists of the free motion of a point particle inside Γ, colliding elastically with the boundary. The motion is then determined by the collision point at Γ and the direction of motion immediately after each impact.

If we parameterize Γ by $\varphi \in \mathbb{R}/2\pi\mathbb{Z}$, the angle between the tangent vector and a fixed direction, the billiard motion can then be described by the parameter φ, that locate the point of impact, and by the angle $\alpha \in (0, \pi)$ between the tangent vector and the outgoing trajectory, measured counterclockwise.

This motion defines a map B from the open cylinder $\mathbb{R}/2\pi\mathbb{Z} \times (0, \pi)$ into itself, which has some very well known properties (see, for instance, [5], chapters 9 and 13 for the properties of billiards and twist maps listed and/or used in this paper): if Γ is C^k, B is a C^{k-1}-diffeomorphism, preserving the measure $d\nu = R(\varphi) \sin\alpha \, d\alpha d\varphi$, where R is the radius of curvature of Γ.

As Γ has strictly positive curvature, B has the monotone Twist property. Therefore, for each $\rho \in (0, 1)$ there exists a closed, invariant, minimal set on the phase space $\mathbb{R}/2\pi\mathbb{Z} \times (0, \pi)$, which can be injectively projected onto $\mathbb{R}/2\pi\mathbb{Z}$ and such that the induced dynamics preserves the order of $\mathbb{R}/2\pi\mathbb{Z}$. This invariant set can be a periodic orbit with period ρ; an invariant *rotational curve*, i.e., a continuous closed curve, non homotopic to a point, invariant by B, with rotation number ρ and which, by Birkhoff's Theorem, is the graph of a Lipschitz function $\alpha = f(\varphi)$; or an *Aubry-Mather Set*, i.e., a closed, invariant, minimal set, projecting injectively on a Cantor set of $\mathbb{R}/2\pi\mathbb{Z}$.

Having an invariant rotational curve is a quite general property: the circular and the elliptical billiards, for instance, are integrable and have the whole cylindrical phase space $\mathbb{R}/2\pi\mathbb{Z} \times (0, \pi)$, for the circle, or a big part of it, for the ellipse, foliated by invariant rotational curves. If the oval boundary Γ is $C^k, k \geq 5$ then Douady-Lazutkin Theorem guarantees a whole family of invariant rotational curves near the boundaries of the cylindrical phase space ([3], [7]).

However only a few ovals will have an invariant horizontal circle. The most famous case is the circle, whose phase space is foliated by invariant horizontal circles given by $\alpha = \alpha_0$. On each one of these circles, the restricted dynamics is a rotation, with rotation number $\omega_0 = \alpha_0/\pi$.

Apart from the circle, there are only two possible cases: ovals of constant width (which includes the circle) or ovals with a very special symmetry (which again includes the circle and some symmetric ovals of constant width).

The first case was studied by Knill [6] who proved that a billiard map preserves the circle $\alpha = \pi/2$ if and only if the oval boundary Γ has constant width. The rotation number of this circle is $1/2$ as each of its points belongs to a 2-periodic orbit.

Tabachnikov [11] and Gutkin [4] proved that a noncircular oval billiard has an horizontal invariant circle of the form $\alpha = \alpha_0 \neq \pi/2$, if and only if the radius of curvature of the boundary Γ is given by $R(\varphi) = a + b\cos(n\varphi)$, with $n \geq 4$ and $|b| < a$, and α_0 satisfies $\tan(n\alpha_0) = n\tan(\alpha_0)$. The billiard dynamics restricted to the invariant horizontal circle is a circle diffeomorphism, with rotation number $\omega_0 = \alpha_0/\pi$ which, as proved by V. Cyr [2], is an irrational number.

In particular, observe that combining these results we have that an oval billiard preserves a horizontal circle with rational rotation number different from a half if and only if its boundary is a circle.

2. Invariant rotational curves near an invariant horizontal circle

In this section we will show that any neighbourhood of an invariant horizontal circle, for sufficiently differentiable billiards, contains an invariant rotational curve. We will separate the proof into three parts corresponding to the three differente cases. We start with the obvious case of the circular billiard. The case of a horizontal circle with rotation number 1/2, is handled by Douady's result ([3], Chapter 3, Corollary II.2). Finally we address the irrational rotation number case. Unfortunately, Douady's result cannot be applied in this last case, since we do not know if the rotation numbers are Diophantine. We will use instead, the ideas developed by Levi and Moser in [8]. Although their result also ask for a Diophantine condition, we will be able to develop a strategy to manage irrational rotations.

2.1. The circular billiard.

If the boundary curve is a circle, then the billiard map is $B(\varphi_0, \alpha_0) = (\varphi_0 + 2\alpha_0, \alpha_0)$. The phase space is foliated by invariant horizontal circles and trivially any neighbourhood of each one of those circles contains other invariant rotational circles.

2.2. Billiards of constant width.

Let Γ be a C^k constant width curve, with length L and width W. The constant width property implies that the billiard map B preserves the horizontal circle $\alpha = \pi/2$ and $B(\varphi, \pi/2) = (\varphi + \pi, \pi/2)$.

Lemma 1. *There is a C^{k-2} change of coordinates on a neighbourhood of the horizontal circle $\alpha = \pi/2$ that puts B into the form*

$$T(t, y) = \left(t + \frac{1}{2} + \zeta(t)y + \mathcal{O}(y^2), y + \mathcal{O}(y^2)\right)$$

where $\zeta_1 \geq \zeta(t) \geq \zeta_2 > 0$.

Proof. Unsing the canonical variables $s \in \mathbb{R}/L\mathbb{Z}$, the arclength parameter of Γ, and $p = \cos\alpha \in (-1,1)$, the billiard map writes $\tilde{B}(s_0,p_0) = (s_1,p_1)$, with $B(\varphi(s_0), \arccos p_0) = (\varphi(s_1), \arccos p_1)$. It is area preserving as the Lebesgue measure $ds\,dp$ is invariant. Clearly, $\tilde{B}(s(\varphi),0) = (s(\varphi+\pi),0)$.

Let $R(\varphi)$ be the radius of curvature of Γ at φ. For $t \in \mathbb{R}/\mathbb{Z}$, the length of Γ from $\varphi = 0$ to $\varphi = 2\pi t$ is given by $s(t) = \int_0^{2\pi t} R(\beta)d\beta$. And so, $\tilde{B}(s(t),0) = (s(t+1/2),0)$. Along with $t = \varphi(s)/2\pi$, we take $y = -2\pi R(\varphi(s))\,p$ to define the change of coordinates

$$G(t,y) = \left(s(t), \frac{-y}{2\pi R(2\pi t)}\right) = \left(s(t), \frac{-y}{s'(t)}\right) \tag{1}$$

In the new variables the billiard map is given by $T = G^{-1} \circ \tilde{B} \circ G$ and we have

$$T(t,0) = G^{-1} \circ \tilde{B}(s(t),0) = G^{-1}(s(t+\frac{1}{2}),0) = (t+\frac{1}{2},0).$$

Observe that $|\det DG_{(t,y)}| = 1$ and so T is also area preserving.

Expanding near $(t,0)$, for each fixed t we get

$$T(t,y) = (t+\frac{1}{2},0) + DT_{(t,0)}(0,y) + \mathcal{O}(y^2).$$

Denoting $W = ||\Gamma(2\pi t) - \Gamma(2\pi t+\pi)||$, $R_0 = R(2\pi t)$, $R_\pi = R(2\pi t+\pi)$, on a constant width curve we have $W = R_0 + R_\pi$. Using the well know expression of the derivative of the billiard map in the (s,p) coordinates (see, for instance [9]) we have

$$\begin{aligned} DT_{(t,0)} &= DG^{-1}_{(s(t+\frac{1}{2}),0)}\, D\tilde{B}_{(s(t),0)}\, DG_{(t,0)} \\ &= \begin{pmatrix} \frac{1}{2\pi R_\pi} & 0 \\ 0 & -2\pi R_\pi \end{pmatrix} \begin{pmatrix} \frac{R_\pi}{R_0} & -W \\ 0 & \frac{R_0}{R_\pi} \end{pmatrix} \begin{pmatrix} 2\pi R_0 & 0 \\ 0 & \frac{-1}{2\pi R_0} \end{pmatrix} = \begin{pmatrix} 1 & \frac{W}{4\pi^2 R_0 R_\pi} \\ 0 & 1 \end{pmatrix} \end{aligned}$$

This leads to the desired expansion

$$T(t,y) = (t+\frac{1}{2},0) + \left(\frac{W}{4\pi^2 R_0 R_\pi}y, y\right) + \mathcal{O}(y^2)$$

for each fixed t.

Since $t \in \mathbb{R}/\mathbb{Z}$ the expansion is uniformly defined on a cylinder $U_\delta = \{(t,y), t \in \mathbb{R}/\mathbb{Z}, |y| < \delta\}$. As Γ is closed, and then compact, and the billiard dynamics is invariant by homotheties on the billiard table, R is positive and bounded above and below, we can assume that $\min R(\varphi) \geq 1/(2\pi)$. Therefore, as W is the (constant) width of the curve, there exist ζ_1 and ζ_2 such that $\zeta_1 \geq W/(4\pi^2 R_0 R_\pi) \geq \zeta_2 > 0$. $\square$

Theorem 2. *Let Γ be a C^k constant width oval, $k > 7$, and B its associated billiard map. Then every cylindrical neighbourhood of the invariant circle $\alpha = \pi/2$ contains an invariant rotational $C^{k'}$-curve, $k' < k-4$, where the billiard dynamics is conjugated to a Diophantine rotation.*

Proof. This theorem is proved applying Douady's result ([3], Chapter 3, Corollary II.2) in the rational $(\alpha/\pi = 1/2)$ case.

The billiard map B is a C^{k-1}-diffeomorphism, preserving the measure $R(\varphi)\sin\alpha\, d\alpha d\varphi$. Changing coordinates (φ,α) to (s,p), $\tilde{B}$ is still a C^{k-1}-diffeomorphism. As it is area preserving, $\tilde{B}^n(\gamma) \cap \gamma \neq \emptyset$ for any rotational curve γ (invariant or not) and any n, i.e., $\tilde{B}$ and all its iterates have the property of intersection. So $T(t, y)$ is a C^{k-2} diffeomorphism which, together with all its iterates, inherits the property of intersection. It can be expanded as $T(t, y) = (t + 1/2 + W[4\pi^2 R(2\pi t)R(2\pi t + \pi)]^{-1} y + \mathcal{O}(y^2), y + \mathcal{O}(y^2))$ by the lemma above. We can then apply Douady's result and conclude that there exists $\epsilon > 0$ and a Cantor set K formed by irrational numbers on $(1/2 - \epsilon, 1/2 + \epsilon)$ such that for every $a \in K$ there is an invariant rotational curve γ_a where the billiard dynamics $B|_{\gamma_a}$ is conjugated to a rotation of $2\pi a$. Moreover, for every $k' < k-4$, $||\gamma_a - \gamma_{1/2}||_{C^{k'}} \to 0$, where $\gamma_{1/2} = \{\alpha = \pi/2\}$. Or, in other words, if (a_n) is a sequence of numbers on $K \setminus \{1/2\}$, $a_n \to 1/2$, the invariant curves γ_{a_n} converge to $\gamma_{1/2}$.

So, every cylindrical neighbourhood of the invariant circle $\alpha = \pi/2$ contains an invariant rotational curve. □



FIGURE 1. Right: Phase space of the billiard on the constant width oval with radius of curvature $R(\varphi) = 1 + 0.1\cos 5\varphi + 0.03\cos 7\varphi$. Left: zoom on a neighbourhood of $\alpha = \pi/2$.

2.3. **Billiards with a special symmetry.**

As we have mentioned in the introduction, Tabachnikov [11] and Gutkin [4] proved that a noncircular oval billiard has an horizontal invariant circle of the form $\alpha = \alpha_0 \neq \pi/2$ if and only if the radius of curvature of the boundary Γ is given by $R(\varphi) = a + b\cos(n\varphi)$, with $n \geq 4$, $|b| < a$, and α_0 satisfies $\tan(n\alpha_0) = n\tan(\alpha_0)$. If n is odd, $\alpha_0 = \pi/2$ is a solution and Γ has constant width. The existence of invariant curves close to this invariant circle is given by Proposition 2 in the previous subsection. In this section we will prove an analogous result for the other circles: every cylindrical neighbourhood of the invariant circle $\alpha = \alpha_0 \neq \pi/2$ contains an analytic and invariant rotational curve, where the billiard dynamics is conjugated to a Diophantine rotation.

As $R(\varphi)$ is analytic, the oval Γ is analytic and so is the billiard map $B(\varphi_0, \alpha_0) = (\varphi_1, \alpha_1)$. Observing that $\varphi_1 - \alpha_1 = \varphi_0 + \alpha_0$ we see that the billiard dynamics restricted to the invariant horizontal circle $\alpha = \alpha_0$, is just a rotation of $2\alpha_0$. However, although Cyr [2] proved that if $\alpha_0 \neq \pi/2$ is a root of $\tan(n\alpha_0) = n\tan(\alpha_0)$ then α_0/π is an irrational number, we do not know if it is Diophantine.

Levi and Moser [8] use the variational formulation of the Twist map to write down a homological equation and an invariant curve is found by a modified Newton's Method. The proof of Levi-Moser's Theorem also strongly relies on the hypothesis that the rotation number is Diophantine and so their result can not be directly applied to our problem. However the steps of the construction can be adapted to handle irrational rotations.

As homotheties on the billiard table do not change the dynamics of the billiard map, we will take Γ with length 1, fixing $a_0 = 1/(2\pi)$. With $|b| < 1/(2\pi)$, Γ is strictly convex and then the billiard problem has a Lagrangian function

$$h(s,\tilde{s}) = -||\Gamma(s) - \Gamma(\tilde{s})||.$$

It is easy to see that (see, for instance, [5], section 9.2)

$$B(s_0,p_0) = (s_1,p_1) \quad \Longleftrightarrow \quad \partial_1 h(s_0,s_1) = p_0 \quad \text{and} \quad \partial_2 h(s_0,s_1) = -p_1$$

A curve $\gamma(t) = (u(t), v(t))$, $t \in \mathbb{R}/\mathbb{Z}$, satisfying

$$B(u(t),v(t)) = (u(t+\omega), v(t+\omega)) \tag{2}$$

is invariant under B and $B|_\gamma$ is conjugated to a rotation of ω. This is the case, for instance, of the invariant horizontal circle $\alpha = \alpha_0$ which corresponds to

$$\begin{aligned} u(t) &= s(t) = \int_0^{2\pi t} R(\beta)\, d\beta = t + \frac{b}{n}\sin 2\pi n t \\ v(t) &= p(t) = \cos\alpha_0 \\ \omega &= \omega_0 = \frac{\alpha_0}{\pi}\ . \end{aligned}$$

Since

$$v(t) = -\partial_1 h(u(t), u(t+\omega)) = \partial_2 h(u(t-\omega), u(t)) \tag{3}$$

formula (2) is true if and only if $u(t)$ satisfies the homological equation

$$\partial_1 h(u(t), u(t+\omega)) + \partial_2 h(u(t-\omega), u(t)) = 0 \tag{4}$$

Instead of looking for an invariant curve we will search for a function u and a number ω verifying (4). In order to do that, we need to set up the framework in which the problem is formulated and solved by given some definitions and preliminary estimates.

Given $r > 0$, let W_r be the space of analytic functions $g(z)$ defined on $|\operatorname{Im} z| \le r$, with the norm $|g|_r = \sup_{|\operatorname{Im} z| \le r} |g(z)|$, and $\hat{W}_r \subset W_r$ the subspace of one-periodic functions.

Lemma 3. *Let $s(z)$ be the analytic continuation of the arclength function $s(t)$. Then $s(z) \in W_r$ and $s(z) - z \in \hat{W}_r$, for every $r > 0$.*

Moreover, given $N > \max\left\{\frac{1}{1-2\pi b}, 1 + 2\pi b\right\}$ there exists $r_0 > 0$ such that $|s'|_r < N$ and $|s'|_r^{-1} < N$ for all $r \le r_0$.

Proof. It is enough to observe that $s(t) = t + (b/n)\sin 2\pi n t$ is analytic. □

Lemma 4. *Let $h(z_1,z_2)$ be the analytic continuation of the Lagrangian function $h(s_1,s_2)$. Its domain D can be chosen such that* $\min|\partial_{12} h(z_1,z_2)| > \kappa$ *and* $|h|_{C^3} < M$ *in D, with $\kappa > 0$ and $M > 0$.*

Proof. $R(\varphi) = 1/2\pi + b\cos n\varphi$ is analytic, and so is the Lagrangian function h. Consider $h(t) = h(s(t), s(t+\omega_0))$, the restriction of the Lagrangian function to the invariant horizontal circle. Then

$$h(t+1) = h(s(t)+1, s(t+\omega_0)+1) = h(s(t), s(t+\omega_0))$$

and h together with all its derivatives up to order three, when evaluated on the invariant horizontal circle, are bounded for every $t \in \mathbb{R}$. Moreover

$$|\partial_{12} h\left(s(t), s(t+\omega_0)\right)| = -\frac{\sin^2\alpha_0}{h\left(s(t), s(t+\omega_0)\right)} \ge C > 0\ .$$

Then we can take an open strip $U \subset \mathbb{R}^2$, containing the curve $(s(t), s(t+\omega_0))$, $t \in \mathbb{R}$, where $|h|_{C^3} < \tilde{M}$ and $|\partial_{12}h(s_1, s_2)| > \tilde{\kappa} > 0$ and extend h, via its Taylor series, to a domain $D \subset \mathbb{C}^2$, $D \cap \mathbb{R}^2 = U$, where $|h|_{C^3} < M$ e $\min_D |\partial_{12}h(z_1, z_2)| > k$. ☐

For $\rho > 0$, let $D_\rho = \{(z_1, z_2) \big| \operatorname{dist}((z_1, z_2), \mathbb{C}^2 - D) > \rho\} \subset D$ be the biggest subset of D such that its ρ-neighbourhood is in D.

Lemma 5. *Given $\rho > 0$ such that $\emptyset \neq D_\rho \subset D$, there exist positive constants r, ξ_ρ and μ_ρ such that for every $|\xi - \omega_0| \leq \xi_\rho$ and z such that $|Imz| \leq r$*

$$(s(z), s(z+\xi)) \in D_\rho$$

and if $|g - s|_r \leq \mu_\rho$ we also have

$$(g(z), g(z+\xi)) \in D_\rho$$

Proof. $\{(s(t), s(t+\omega_0)), t \in \mathbb{R}\} \subset D$. The continuity of s and the equality $s(z+1) = s(z) + 1$ imply the first result. The second one is a consequence of the first. ☐

Given $\rho > 0$, let

$$U_{\mu_\rho} = \left\{ g \in \hat{W}_r \middle| \; |g - s|_r \leq \mu_\rho \right\}.$$

For each $\xi \in [\omega_0 - \xi_\rho, \omega_0 + \xi_\rho]$ let

$$E_\xi : U_{\mu_\rho} \longrightarrow \hat{W}_r,$$
$$E_\xi(g)(z) = \partial_1 h(g(z), g(z+\xi)) + \partial_2 h(g(z-\xi), g(z)) \;.$$

Levi and Moser in [8] proved

Theorem (Levi-Moser) *Suppose that ξ verifies, for all $n, m \in \mathbb{Z}$, $n > 0$, a Diophantine condition $|n\xi - m| \geq K/n^{\beta+1}$. Then there exist positive numbers δ, r_∞ and A, depending on $\rho, r, \kappa, M, N, \beta, K$ such that if $|E_\xi(u)|_r < \delta$ then there exists a unique u_ξ such that $E_\xi(u_\xi) = 0$, $u_\xi(\theta) - \theta \in \hat{W}_{r_\infty}$ and $|u_\xi - u|_{r/2} < A|E_\xi(u)|_r$.*

Obviously $E_{\omega_0}(s) = 0$; however, as we already have pointed out, we do not know if ω_0 is a Diophantine number. In the next steps we show how to circumvent this obstacle.

Lemma 6. *Given $\delta_1 > 0$ there exists $\epsilon_1 > 0$ such that for every $|\xi - \omega_0| < \epsilon_1$, $|E_\xi(s)|_r < \delta_1$.*

Proof. Let $f : \{|\operatorname{Im} z| \leq r\} \times [\omega - \xi_\rho, \omega + \xi_\rho] \to \mathbb{C}$ be the uniformly continuous function given by

$$f(z, \xi) = \partial_1 h(s(z), s(z+\xi)) + \partial_2 h(s(z-\xi), s(z)).$$

Then given $\delta_1 > 0$, there exists $\epsilon_1 > 0$ such that for all $|(z_1, \xi) - (z, \omega_0)| < \epsilon_1$, we have $|f(z_1, \xi) - f(z, \omega_0)| < \delta_1/2$.

Since $f(z, \omega_0) = 0$ we can take $z_1 = z$ to conclude that given $\delta_1 > 0$ there exists $\epsilon_1 > 0$ such that if $|\xi - \omega_0| < \epsilon_1$ then $|f(z, \xi)| = |E_\xi(s(z))| < \delta_1/2 < \delta_1$. ☐

Lemma 7. *Let $\lambda \in (0,1)$ and $\varepsilon > 0$ be two numbers such that $[\lambda - \varepsilon, \lambda + \varepsilon] \subset [0,1]$ and let $K_0 = \min\{\lambda - \varepsilon, 1 - (\lambda + \varepsilon)\}$. For each fixed K_1 and β_1, $K_0 \geq K_1 > 0, \beta_1 > 0$, the set*

$$C_{\lambda,\varepsilon}(K_1, \beta_1) = \{\xi \in [\lambda - \varepsilon, \lambda + \varepsilon] \text{ such that } |n\xi - m| \geq \frac{K_1}{n^{2+\beta_1}}, \forall m, n \in \mathbb{Z}, n > 0\}$$

has full Lebesgue measure in $[\lambda - \varepsilon, \lambda + \varepsilon]$.

Proof. If $\xi \in C_{\lambda,\varepsilon}(K_1,\beta_1)$ we have in fact $|n\xi - m| \geq \dfrac{K_1}{n^{2+\beta_1}}$ for all m and $n>1$.

Let us denote $I_{\lambda,\varepsilon} = [\lambda-\varepsilon, \lambda+\varepsilon]$ and define the sets

$$A(\beta_1) = \left\{ \xi \in I_{\lambda,\varepsilon} \,\middle|\, |n\xi - m| \geq \frac{k}{n^{2+\beta_1}} \ \forall m, \forall n > 1, \text{ and at least one } 0<k\leq K_0 \right\}$$

$$B(k,\beta_1) = \left\{ \xi \in I_{\lambda,\varepsilon} \,\middle|\, |n\xi - m| < \frac{k}{n^{2+\beta_1}} \ \text{ for some } m \text{ and } n>1 \right\} .$$

It is clear that $A \supset C_{\lambda,\varepsilon}(K_1,\beta_1)$ for all $0<K_1\leq K_0$. Note that the complement of $A(\beta_1)$ in $I_{\lambda,\varepsilon}$ is the set $A^c(\beta_1) = \bigcap\limits_{k\in(0,K_0]} B(k,\beta_1)$.

For each fixed m and n the set of ξ such that $|n\xi - m| < \dfrac{k}{n^{2+\beta_1}}$ is the interval $I(k,\beta_1,n,m)$ of lenght $\dfrac{2k}{n^{3+\beta_1}}$. On the other hand, as the inequality also implies that $-1<m<n+1$, it follows that

$$B(k,\beta_1) \subset \bigcup_{n=2}^{\infty}\left(\bigcup_{m=0}^{n} I(k,\beta_1,n,m)\right)$$

and so, if μ denotes the Lebesgue measure,

$$\mu\left(B(k,\beta_1)\right) \leq \sum_{n=2}^{\infty}\left(\sum_{m=0}^{n}\mu\left(I(k,\beta_1,n,m)\right)\right) \leq 3k\sum_{n=2}^{\infty}\frac{1}{n^{2+\beta_1}} \ , \ \forall k>0 \ .$$

Thus $\mu\left(A^c(\beta_1)\right) \leq kC$ for all k, which implies that $\mu\left(A^c(\beta_1)\right)=0$.

Now, for each fixed $K_1 \leq K_0$, let us define the set

$$D(K_1) = \left\{ \xi \in I_{\lambda,\varepsilon} \,\middle|\, |n\xi - m| \geq \frac{K_1}{n^{2+\beta}}, \ \forall m, \forall n>1, \text{ and at least one } \beta>0 \right\} .$$

Clearly $D(K_1) \supset C_{\lambda,\varepsilon}(K_1,\beta_1)$ and $D^c(K_1) = \bigcup_{\beta} B(K_1,\beta)$. A similar argument shows that $\mu\left(D^c(K_1)\right)=0$. As $C_{\lambda,\varepsilon}(K_1,\beta_1) = A(\beta_1)\cap D(K_1)$, $\mu(C^c_{\lambda,\varepsilon}(K_1,\beta_1)=0$ and so $\mu(C_{\lambda,\varepsilon}(K_1,\beta_1)) = \mu(I_{\lambda,\varepsilon}) = 2\varepsilon$. ☐

We can now prove the main result of this subsection.

Theorem 8. *Let Γ be an oval such that its associated billiard map B preserves the horizontal circle given by $\alpha=\alpha_0\neq\pi/2$. Then every cylindrical neighbourhood of the invariant circle contains an analytic and invariant rotational curve, where the billiard dynamics is conjugated to a Diophantine rotation.*

Proof. By Levi-Moser's Theorem if $\xi \in I_{\omega_0,\xi_\rho} = [\omega_0-\xi_\rho, \omega_0+\xi_\rho]$ is a Diophantine number with constants β and K, there exist δ and r_∞ such that if $\left|E_\xi\left(u\right)\right|_r < \delta$ then there exists a unique u_ξ such that $E_\xi\left(u_\xi\right)=0$ where $u_\xi\left(z\right)-z\in\hat{W}_{r_\infty}$, the numbers δ and r_∞ depend on $\rho, r, \kappa, M, N, \beta$ and K.

Yet, for any Diophantine number $\xi \in I_{\omega_0,\xi_\rho}$ the constants ρ, r, κ, M, N are always the same since, as the oval is fixed, we are dealing with the same Lagrangian function h and the same invariant horizontal circle, given by $u=s$.

Choosing $\varepsilon<\xi_\rho$, $\beta_1>0$ and K_1 as in Lemma 7, all $\xi\in C_{\omega_0,\varepsilon}(K_1,\beta_1)\subset I_{\omega_0,\xi_\rho}$, share the constants K_1 and β_1 so the same δ and r_∞ of Levi-Moser's Theorem works for all of them. Let $\delta_1<\delta=\delta(\rho,r,\kappa,M,N,\beta_1+1,K_1)$. Then, by Lemma 6, there exists $\epsilon_1<\xi_\rho$ such that for every $\xi\in C_{\omega_0,\varepsilon}(K_1,\beta_1)\cap I_{\omega_0,\epsilon_1}$, we have $E_\xi(s)<\delta_1<\delta$.

Thus, for each such ξ, there exists a unique u_ξ such that $E_\xi\left(u_\xi\right)=0$ and so there is an invariant and analytic rotational curve $(u_\xi(t), v_\xi(t))$, where v is given by formula (3), on a neighbourhood of the invariant horizontal circle $(u(t), v(t))=(s(t), \cos\alpha_0)$ and with rotation number ξ.

Now, let $U_\eta=\left\{(s,p)\in\mathbb{R}^2/\, |p-\cos\alpha_0|<\eta\right\}$ be an open strip around the invariant horizontal circle with width 2η. The Lagrangian function h is analytic and verifies $h(s_1+1, s_2+1)=h(s_1,s_2)$ and so $\partial_1 h$ is uniformly continuous in $\mathbb{R}^2$. So, there exists $\nu>0$ such that if $|(\sigma_1,\sigma_2)-(s_1,s_2)|<\nu$ then $|\partial_1 h(\sigma_1,\sigma_2)-\partial_1 h(s_1,s_2)|<\eta$. Levi-Moser's Theorem states that the solution u_ξ of $E_\xi=0$ satisfies $|u_\xi-s|_{r/2}<A|E_\xi(s)|_r$ for some positive constant A. So for $\xi\in C_{\omega_0,\varepsilon}(K_1,\beta_1)$, taking $|\xi-\omega_0|$ sufficiently small, we can have, by lemma 6, $|E_\xi(s)|_r<\nu/(\sqrt{2}A)$, obtaining $|u_\xi-s|_{r/2}<\nu/(\sqrt{2})$ and then

$$|(u_\xi(z), u_\xi\left(z+\xi\right))-\left(s\left(z\right), s\left(z+\xi\right)\right)|<\nu.$$

Then, for $t\in\mathbb{R}$

$$|v_\xi(t)-\cos\alpha_0|=|\partial_1 h(u_\xi(t), u_\xi(t+\xi))-\partial_1 h(s(t), s(t+\xi))|<\eta$$

and $\gamma_\xi=\{(u_\xi(t), v_\xi(t))\}\subset U_\eta$ or, in words, any cylindrical neighbourhood of the invariant horizontal circle contains an analytic and invariant rotational curve, where the dynamics is conjugated to a Diophantine rotation. □



FIGURE 2. Right: Phase space of the billiard on the symmetric oval with radius of curvature $R(\varphi)=1+0.1\cos 6\varphi$. Left: zoom on a neighbourhhod of the horizontal circle $\alpha=\alpha_0\approx 0.75625271$.

3. DYNAMICAL CONSEQUENCES

Birkhoff in [1] proved that any rotational invariant curve is a Lipschitz graph $\alpha=f(\varphi)$. We can define a partial order on the set of invariant rotational curves: given two invariant rotational curves $\gamma_1=\{(\varphi, f_1(\varphi))\}$ and $\gamma_2=\{(\varphi, f_2(\varphi))\}$, we say that γ_1 is bellow γ_2, denoted by $\gamma_1<\gamma_2$, if $f_1(\varphi)<f_2(\varphi)$. If $f_1(\varphi)>f_2(\varphi)$ we say that γ_1 is above γ_2.

Theorem 9. *Let B be a billiard map preserving a horizontal circle $\gamma_0=\{\alpha=\alpha_0\}$, defined inside a C^k-curve, $k>7$, if $\alpha_0=\pi/2$, or inside an analytic curve otherwise. Then γ_0 can be approached by invariant rotational curves from above and from bellow.*

Proof. We already know the existence of rotational invariant curves approaching the invariant horizontal circle. The question here is to prove that they can not approach the horizontal circle only by one side.

The result is obvious for the circular billiard.

For the case $k > 7$ and $\alpha_0 = \pi/2$ it is sufficient to remark that the billiard map is reversible with respect to the reversing symmetry $S(\varphi, \alpha) = (\varphi, \pi - \alpha)$, as $S \circ B = B^{-1} \circ S$, meaning that the phase space is symmetric with respect to the line $\alpha = \pi/2$.

If $\alpha_0 \neq \pi/2$, i.e. $\omega_0 \neq 1/2$, for every $\xi \in C_{\omega_0,\, \varepsilon}(K_1, \beta_1,) \cap (\omega_0 - \epsilon_1, \omega_0 + \epsilon_1)$ there exists a unique invariant and analytic rotational curve $\gamma_\xi = \{(u_\xi(t), v_\xi(t))\}$, with rotation number ξ.

Moreover, the Lebesgue measure of $C_{\omega_0,\, \varepsilon}(K_1, \beta_1) \cap (\omega_0 - \epsilon_1, \omega_0 + \epsilon_1)$ is $2\epsilon_1$. Then there are invariant curves with rotation number $\xi < \omega_0$ and also with $\xi > \omega_0$. It is proved in [10] (Corollary 6, page 110) for general Twist maps, that if two invariant rotational curves are such that γ_1 is bellow γ_2, then the rotation number of γ_1 is smaller then the rotation number of γ_2. This implies that all the invariant rotational curves given by Proposition 8 can not be on the same side of the horizontal circle, since they would have rotation numbers always greater or always smaller than ω_0. □

In the same work [1], Birkhoff defined an instability region as the region between two invariant rotational curves, with no other invariant rotational curves inside. An immediate consequence of Theorem 9 is

Corollary 10. *On sufficiently differentiable oval billiards, no instability region has an horizontal circle as a boundary.*

The orbit of a point (φ_0, α_0), the set $O(\varphi_0, \alpha_0) = \{(\varphi_n, \alpha_n) = B^n(\varphi_0, \alpha_0)\}$, is ordered if $\Gamma(\varphi_0) < \Gamma(\varphi_1) < \Gamma(\varphi_2)$ implies $\Gamma(\varphi_n) < \Gamma(\varphi_{n+1}) < \Gamma(\varphi_{n+2})$, for every n, in the order induced by the orientation of the boundary Γ.

Lemma 11. *For sufficiently differentiable oval billiards having an invariant horizontal circle γ_0, there is an invariant region $\mathcal{R}$, containing γ_0, where every orbit is ordered.*

Proof. Using the expression of derivative of the billiard map in the (φ, α) coordinates, the Taylor expansion of B is given by

$$B(\varphi, \alpha_0 + \alpha) = (\varphi + 2\alpha_0, \alpha_0) + \frac{\alpha}{\sin \alpha_0} \left(L, \frac{L}{R(\varphi + 2\alpha_0)} - \sin \alpha_0 \right) + \mathcal{O}(\alpha^2)$$

near $\gamma_0 = \{\alpha = \alpha_0\}$. As Γ is closed, this implies that there is an ϵ sucht that if $|\alpha| < \epsilon$ then the orbit $O(\varphi, \alpha_0 + \alpha)$ is ordered. Let $\gamma_1 < \gamma_0 < \gamma_2$ be two invariant rotational curves contained in $\{(\varphi, \alpha_0 + \alpha), |\alpha| < \epsilon\}$ and $\mathcal{R}$ be the region between γ_1 and γ_2. $\mathcal{R}$ is invariant, contains γ_0 and every orbit in it is ordered. □

Corollary 12. *For a sufficiently differentiable oval billiard, the set of invariant curves has positive measure on $\mathcal{R}$.*

Proof. If $\alpha_0 = \pi/2$ and the oval is C^k for k sufficiently large, by Douady's observation ([3],chapter 3, II.3.3) the Cantor set K obtained in the proof of Theorem 2 can be replaced by the set

$$K\left(\beta, \varepsilon\right) = \left\{ a \in \mathbb{R} \text{ such that } \left| a - \frac{1}{2} \right| \leq \varepsilon \text{ and for all } \left| a - \frac{m}{n} \right| \geq \lambda \frac{\left| a - \frac{1}{2} \right|}{n^{2+\beta}} \right\}$$

where $\beta > 0$ satisfies $7 + 2\beta < k$ and $0 < \varepsilon < \varepsilon_0$ wich ε_0 depending on $\lambda > 0$ and $\beta > 0$. He also notes that $\lim_{\varepsilon \to 0} \mu\left(K\left(\beta, \varepsilon\right)\right)/\varepsilon = 1$ and this implies that $\mu\left(K\left(\beta, \varepsilon\right)\right) > 0$.

Let γ_1 and γ_2 be the boundaries of the invariant region $\mathcal{R}$ of Corollary 11 and ξ_1 and ξ_2 be their respectively rotation numbers. We can assume $K(\beta, \varepsilon) \subset [\xi_1, \xi_2]$.

By proposition 13.2.7 from Katok and Hasselblat's book ([5], page 427), the rotation number function $\rho : \mathcal{R} \mapsto [\xi_1, \xi_2]$ is continuous in the topology of the Hausdorff metric.

Once that $\mu(K(\beta, \varepsilon)) > 0$ and $K(\beta, \varepsilon) \subset [\xi_1, \xi_2]$ then $\rho^{-1}([\xi_1, \xi_2])$ has positive measure in $\mathcal{R}$.

We can repeat the argument in the case $\alpha_0 \neq \pi/2$. Lemma 7 assures that the set of Diophantine numbers corresponding to invariant rotational curves on $\mathcal{R}$ has full Lebesgue measure on $[\xi_1, \xi_2]$. Then $\rho^{-1}([\xi_1, \xi_2])$ has positive measure in $\mathcal{R}$. $\square$

Acknowledgements: This work was partially supported by the Brazilian agencies CNPq and FAPEMIG.

References

[1] G.D. Birkhoff: *Sur quelques courbes fermées remarquables.* Bull. SMF **60** (1932) 1–26. Also in *Collected Math. Papers of G. D. Birkhoff* **II**, 444–461.

[2] V. Cyr, *A Number Theoretic Question Arinsing in the Geometry of Plane Curves and in Billiard Dynamics*, arXiv:1103.5071v1, 2011.

[3] R. Douady: *Applications du théorème des tores invariants.* Thèse de 3ème Cycle, Univ. Paris VII, 1982.

[4] E. Gutkin: *Capillary floating and the billiard ball problem.* Preprint arXiv:1012.2448, 2010.

[5] A. Katok, B. Hasselblatt: *Introduction to the Modern Theory of Dynamical Systems.* Univ. Cambridge Press, 1997.

[6] O. Knill: *On nonconvex caustics of convex billiards.* Elem. Math. **53** (1998), 89–106.

[7] V.F. Lazutkin: *The existence of caustics for a billiard problem in a convex domain.* Math. USSR Izvestija **7** (1973), 185–214.

[8] M. Levi, J. Moser: *A Lagrangian Proof of the Invariant Curve Theorem for Twist Mappings,* Proceedings of Symposia in Pure Mathematics, volume 69, AMS, 733-746, 2001.

[9] S. Oliffson Kamphorst, S. Pinto-de-Carvalho: *The first Birkhoff coefficient and the stability of 2-periodic orbits on billiards*, Exp.Math, **14/3**, 299-306, 2005.

[10] C.G. Ragazzo, M. J. Dias Carneiro, S. A. Zanata: *Introdução à Dinâmica de Aplicações do Tipo Twist*, Publ.Mat., IMPA, 2005.

[11] S. Tabachnikov: *Billiards*, Société Mathématique de France, 1995.

Dep. de Matemática, ICEB, UFOP, Ouro Preto, Brasil
E-mail address: `gegpreto@gmail.com`

Dep. de Matemática, ICEx, UFMG, Belo Horizonte, Brasil
E-mail address: `syok@mat.ufmg.br`

Dep. de Matemática, ICEx, UFMG, Belo Horizonte, Brasil
E-mail address: `sonia@mat.ufmg.br`